



COMMISSION  
**OSPAR**

# Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

## Évaluation de l'Indicateur Commun



# OSPAR

## BILAN DE SANTÉ 2023

## 2023

# Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

## OSPAR Convention

The Convention for the Protection of the Marine Environment of the North-East Atlantic (the "OSPAR Convention") was opened for signature at the Ministerial Meeting of the former Oslo and Paris Commissions in Paris on 22 September 1992. The Convention entered into force on 25 March 1998. The Contracting Parties are Belgium, Denmark, the European Union, Finland, France, Germany, Iceland, Ireland, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom.

## Convention OSPAR

La Convention pour la protection du milieu marin de l'Atlantique du Nord-Est, dite Convention OSPAR, a été ouverte à la signature à la réunion ministérielle des anciennes Commissions d'Oslo et de Paris, à Paris le 22 septembre 1992. La Convention est entrée en vigueur le 25 mars 1998. Les Parties contractantes sont l'Allemagne, la Belgique, le Danemark, l'Espagne, la Finlande, la France, l'Irlande, l'Islande, le Luxembourg, la Norvège, les Pays-Bas, le Portugal, le Royaume-Uni de Grande Bretagne et d'Irlande du Nord, la Suède, la Suisse et l'Union européenne

## Contributeurs

Auteurs principaux : Arnaud Louchart, Matthew Holland, Abigail McQuatters-Gollop and Luis Felipe Artigas

Avec le soutien des groupes suivants : OSPAR Pelagic Habitat Expert group, Intersessional Correspondence Group on the Coordination of Biodiversity Assessment and Monitoring, Biodiversity Committee and the OSPAR Secretariat.

## Réalisation de l'évaluation

Ce travail a été co-financé par le Fonds européen pour les affaires maritimes et la pêche dans le cadre du projet : « North-East Atlantic project on biodiversity and eutrophication assessment integration and creation of effective measures (NEA PANACEA) », financé par la DG ENV/MSFD 2020 de l'Union européenne, dans le cadre de l'accord n° 110661/2020/839628/SUB/ENV.C.2.



Co-funded by  
the European Union

## Citation

Louchart, A., Holland, M., McQuatters-Gollop, A. and Artigas, L. F. 2023. *Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton*. OSPAR, 2023: Bilan de santé de l'Atlantique du Nord-Est. Commission OSPAR, Londres. Disponible via le lien suivant : <https://oap.ospar.org/fr/ospar-assessments/quality-status-reports/qsr-2023/evaluations-des-indicateurs/modifications-biomasse-phytoplancton-abondance/?edit&language=fr>

## Table des matières

Contributeurs .....	1
Citation .....	1
Message clé .....	3
Contexte .....	3
Contexte (version étendue).....	3
Méthode d'évaluation .....	4
Résultats .....	18
Résultats (version étendue).....	21
Conclusion .....	26
Conclusion (version étendue).....	26
Lacunes dans les connaissances .....	26
Lacunes dans les connaissances (version étendue) .....	26
Références .....	28
Métadonnées d'évaluation .....	31

## Message clé

La biomasse du phytoplancton et l'abondance du zooplancton ont changé de manière significative (2015-2019), très probablement sous l'effet du changement climatique et des nutriments. Les deux composantes ont diminué dans les habitats océaniques (golfe de Gascogne), ainsi que dans les habitats à salinité variable et les habitats de plateau (mer du Nord au sens large, mers celtiques). Le plancton a connu des tendances contrastées dans tous les habitats côtiers et les habitats du plateau du golfe de Gascogne et de la côte ibérique.

## Contexte

Les algues et les animaux microscopiques à la dérive constituent la base du réseau trophique pélagique et sont une source de nourriture directe ou indirecte pour les poissons, les crustacés et les oiseaux de mer. Les organismes planctoniques sont très sensibles aux facteurs physiques et chimiques, notamment les teneurs en nutriments, la salinité et la température. Ces facteurs dépendent des variations naturelles du climat et de l'hydrographie, ainsi que des processus anthropiques. Les communautés planctoniques ont un cycle de vie court et elles réagissent donc plus rapidement que des niveaux trophiques supérieurs aux changements environnementaux. Les indicateurs se fondant sur le plancton peuvent aider à déceler les modifications de la quantité du plancton et de la structure des communautés planctoniques, qui peuvent avoir un impact sur la structure et le fonctionnement des écosystèmes marins.

L'indicateur 2 relatif aux habitats pélagiques (PH2) concerne les « modifications de la biomasse du phytoplancton et de l'abondance du zooplancton ». Cet indicateur permet d'identifier les modifications (anomalies) dans les quantités de deux groupes fondamentaux au sein d'une communauté planctonique, le phytoplancton et le zooplancton (évalué par l'abondance des copépodes, qui constituent le groupe zooplanctonique le plus nombreux). Ces modifications représentent des déviations de la variabilité naturelle présumée dans les séries temporelles sur le plancton. Les modifications de la biomasse du phytoplancton et de l'abondance du zooplancton sont mesurées entre une période de comparaison passée (avant 2015) et une période d'évaluation temporaire (2015 - 2019). La direction du changement est statistiquement identifiée comme étant soit en augmentation, soit inchangée, soit en diminution. Cet indicateur a été évalué à l'échelle régionale, en utilisant les unités d'évaluation COMP4 (Enserink et al., 2019) pour diviser les données fournies pour la mer du Nord au sens large, les Mers celtiques et le golfe de Gascogne et la côte ibérique (Régions OSPAR II, III et IV). L'assimilation des résultats de cet indicateur avec les changements dans les formes de vie du plancton et la diversité du plancton aidera à fournir une compréhension plus holistique des changements qui se produisent dans l'ensemble des habitats pélagiques.

## Contexte (version étendue)

The quantity of marine plankton is, to a large extent, determined by nutrient concentrations, climate, hydrodynamic drivers and food availability (e.g., Beaugrand *et al.*, 2009). Total phytoplankton (as biomass using chlorophyll-*a* or Phytoplankton Colour Index as a proxy) and zooplankton (as abundance - using total copepods abundance) represent key components of the plankton community. They account for the largest part of the plankton biomass and thus play an important role in overall plankton production, as well as grazing, lysis (auto- and viral), advection and sedimentation processes. Being at the base of the food web, plankton represent (directly or indirectly) a food resource for numerous species at higher trophic levels, such as fish of commercial interest (mackerel, herring, cod). Variability in phytoplankton biomass and zooplankton abundance can have significant impacts on the structure and function of the marine food web, as well as on ecosystem processes such as nutrient recycling. The intrinsic characteristics of plankton organisms, such as their small size, lack of commercial exploitation, short lifecycles, and a global distribution, make them essential components of monitoring programmes aiming to assess the state of marine ecosystems. Their short lifecycles and sensitivity to abiotic factors mean that plankton indicators can provide early warnings of

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

change in the marine ecosystem (Batt *et al.*, 2013), from both natural and human-induced change, over a wide range of temporal and spatial scales. Increased pressure on marine systems is expected to lead to greater and more frequent changes at the base of the food web. Early warnings of ecosystem change can help to prompt management actions when changes are still manageable (see Burthe *et al.*, 2016).

Robust statistical techniques exist to identify significant components of variation and changes at multiple temporal and spatial scales for plankton. These changes may indicate major changes in the marine system involving consequences for other ecosystem components and processes. Significant changes in this indicator are evaluated through time-series analysis. One advantage of the methodology is its ease of application to any data set that can be considered 'long enough' to represent the temporal variability of plankton. However, since plankton dynamics are not well understood, it is difficult to decide the minimum appropriate length of a time-series for an assessment. As a starting point, it is recommended that the minimum length of a time-series should be five years, but preferably a minimum of ten years. It is also necessary that monitoring data are acquired at regular intervals through consistent sampling and analytical procedures.

The methodology for this indicator can be applied to fixed monitoring station time-series (the most common coastal monitoring in European countries) and to large-scale spatio-temporal data sets such as the Continuous Plankton Recorder (CPR) or more recently satellite data. The three data types are complementary and may provide information on different temporal / spatial scales and pressures in the future. For instance, coastal data from fixed monitoring stations can be used to identify plankton indicators that could be linked to human pressures, while large-scale spatio-temporal data in the open ocean can be used to identify plankton indicators that could be linked to large-scale hydro-meteorological changes or to indirect human pressures (e.g., fishing). An important advantage of these plankton indicators is that the concepts are relatively easily transferable to other regions (Gowen *et al.*, 2011; Rombouts *et al.*, 2013). For the future development of these indicators the definition of comparison periods will require knowledge of environmental and human pressure data.

The OSPAR Quality Status Report 2010 highlighted the potential impacts of climate change and other human pressures on plankton communities. Phytoplankton chlorophyll-*a* and phytoplankton indicator species are also assessed under the [Common Procedure for the Identification of the Eutrophication Status of the OSPAR Maritime Area](#) (OSPAR Agreement 2022-07) but there was no comparable regional assessment of phytoplankton biomass and zooplankton abundance due to the use of different assessment units divisions.

A current challenge is to separate expected natural variability from the variability induced by human pressures, a particularly difficult objective that is not yet resolved by plankton science. In the future, further development of this indicator will focus on making the link with human pressures and environmental / climate variability (see Buttay *et al.*, 2015).

### Méthode d'évaluation

The full method employed for this indicator is further elaborated in an OSPAR guidance document for both phytoplankton and zooplankton (Aubert *et al.*, 2017, Duflos *et al.*, 2018, OSPAR PH2 CEMP guidelines, OSPAR Agreement 2019-06).

The methodology has been adapted since the Intermediate Assessment (IA) 2017 (mainly in the first steps of data preparation) due to the type of data used. First, phytoplankton and zooplankton are considered separately. Second, three main data types related to different acquisition systems are considered: time-series of plankton samples collected at fixed (mainly coastal) stations, plankton data from semi-autonomous collecting devices that regularly cover large spatial domains, such as the Continuous Plankton Recorder (CPR) data set, and satellite chlorophyll-*a* (ocean colour product) that provide a monthly synoptic view of the OSPAR Regions.

### Pre-analysis steps: Specificities related to data type

### *Phytoplankton data*

Instead of querying a particular species or group, the bulk phytoplankton community is examined through the total phytoplankton biomass. Phytoplankton biomass can be measured as biovolume, carbon content, or can be assessed through a proxy by measuring the concentration of chlorophyll-*a*, a pigment which is present in all phytoplankton organisms. It is also possible to derive a semi-quantitative measurement of phytoplankton biomass by analysing the Phytoplankton Colour Index (PCI), a metric of pigmentation specific to CPR data. This method estimates the green colour of the plankton community sampled onto a silk net (McQuatters-Gollop *et al.*, 2015). Despite their shorter time-series (since 1997 with SeaWiFS derived data), satellite data were preferable for this analysis over PCI as they provide a more integrated and synoptic view with more regular monitoring and fine-scale spatial and temporal resolution which facilitated comparisons across all areas, including those not monitored by CPR. *In-situ* chlorophyll-*a* and satellite chlorophyll-*a* are used in this assessment as they represent the two types of data most regularly distributed in all areas (including reference stations and ocean colour data from remote sensing algorithms). For data from discrete stations no spatial aggregation was required prior to time series analysis. For non-station data (e.g., satellite, CPR), see the below section “*Non-station Data*”.

### *Zooplankton data*

For zooplankton, only total copepod abundance was included in the indicator calculation. Expression of total zooplankton abundance as total copepod abundance was justified, given that copepods are readily identifiable in samples and represent the balance between production/import and mortality/export between phytoplankton and zooplankton (Budria *et al.*, 2017). In addition, copepods are generally the most abundant and ubiquitous zooplankton taxa, both in space and time (e.g. Rombouts *et al.*, 2009, Turner, 2004, Ikeda, 1985). In practice, the use of groups, such as copepods, is often favoured over single species as indicators (De Jonge, 2007). Indeed, some species, such as those belonging to meroplankton, can exhibit very patchy distribution and highly variable fluctuation in abundance between years. These fluctuations are often due to natural physical dynamics rather than human pressures (De Jonge, 2007). An indicator based on only one species is also unlikely to represent the whole trophic level of zooplankton communities, and which is required here for the present indicator assessment. Using a taxonomic group as broad as copepods we represent a variety of trophic interactions, characteristic of the zooplankton community, which will enable us to detect changes in the pelagic habitat. For data from discrete stations no spatial aggregation was required prior to time series analysis. For non-station data (e.g., satellite and CPR), see the “*Non-station Data*” section.

### *Non-station data*

Large spatio-temporal pelagic data sets originated from satellite for phytoplankton biomass and from the CPR for zooplankton abundance. The satellite data provided by Plymouth Marine Laboratory (PML) originated from the European Space Agency Ocean Colour Climate Change Initiative using the OC5CI Chla algorithm. The OC5CI chlorophyll algorithm can be applied in case 1 - open ocean regions, as well as case 2 - coastal regions (Tilstone *et al.*, 2022). The satellite provided by Royal Belgian Institute of Natural Sciences (RBINS) originated from the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) using the CHL-Gons algorithm from Sentinel3/OLCI. This algorithm is developed for turbid and eutrophic waters. Prior to analysis all copepod species abundance per unit of time (within a sample) were summed to obtain total copepods abundance. The first common step of both plankton components was to aggregate data in space and time using a 1-degree grid for each month to better assess the study area. The procedure consisted of rounding latitude and longitude of samples to the nearest degree. If several sites within a pixel of the grid were acquired during the same month, the average value was calculated. The second step consisted in computing temporal interpolation of the pixels of the grid. For each cell of the grid, if more than 3 consecutive months or more than 4 non-consecutive months were missing in a year, the whole year was removed from the time-series. Thus, for time-series with 3 consecutive or 4 non-consecutive missing months within a year, temporal interpolation was processed. The temporal interpolation was particularly useful in supplementing

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

the year-round satellite data for the North Sea, which are often missing during the winter months due to the short daylight duration (November to February).

For assessment at large spatial scale, chlorophyll-*a* estimation from satellite (both PML and RBINS) as well as zooplankton abundance from the Continuous Plankton Recorder (CPR, Marine Biological Association) are preferred over *in situ* data originated from monitoring fixed stations.

### **Methodology and concept:**

This indicator is based on identification of phytoplankton biomass and zooplankton abundance anomalies within plankton time-series. Anomalies represent deviations from the assumed natural variability of a time-series. Thus, the greater the magnitude of the anomaly (in terms of absolute value, since anomalies can be positive or negative), the greater the change. An anomaly value of zero indicates no difference from the time-series mean (which must be de-seasonalised). To understand the changes presented (i.e., annual anomalies) and to be most useful for decision makers, annual anomalies are best interpreted with information provided by anomalies on monthly time-scales. An R-script for the plankton time series was first developed by Ibanez (reported in Berline *et al.*, 2009), and then adapted for this assessment.

### **Previous assessment:**

The previous assessment (OSPAR Intermediate Assessment; IA 2017) was based on the establishment and the categorisation of annual anomalies of plankton biomass. Anomalies were categorised to improve the presentation of results from a graphical perspective and to simplify the results for management use. The categorisation is based on percentiles using the 2,5, 25, 50, 75 and 97,5<sup>th</sup> percentiles for each time-series. Three categories were used: *small change* (anomalies within the 25–75<sup>th</sup> percentile range), *important change* (anomalies within the 2,5–25<sup>th</sup> percentile range and 75–97,5<sup>th</sup> percentile range) and *extreme change* (anomalies within the 0–2,5<sup>th</sup> percentile range and 97,5–100<sup>th</sup> percentile range). Anomalies within the *small change* category represent the scenario of least likely occurring significant shifts at the plankton community level, and thus the least likely to impact the marine ecosystem. Anomalies within the *important change* and *extreme change* categories have increasing potential to represent significant modifications in the plankton community and the marine ecosystem. The present assessment is based on the previous assessment methodology with some improvements since IA2017. Most of these improvements have already been described and applied in the French national MSFD assessment (Duflos *et al.*, 2018).

### **Temporal trend analysis of biomass anomalies**

Temporal resolution increased between IA2017 and the present assessment from yearly anomalies to monthly calculation. This change in methodology was necessary to better capture variations in phytoplankton biomass and zooplankton abundance beyond their natural cycles. This increase of temporal resolution could be achieved through monthly or higher sampling frequency.

When the data are in the format of monthly mean values, they can be fitted to the COMP4 assessment units (see the subsection **Spatial scales of the assessment**). Following these steps, the time-series analysis can be run. Both phytoplankton and zooplankton time-series analyses are run using the same R-script for both discrete-station data and non-station data, after the pre-analysis steps have been followed. The first step consists of identifying the mean seasonal cycle (which is called seasonality in this assessment) during the whole study period. Removing the seasonality is required in order to analyse the variations of each plankton compartment (i.e., phytoplankton biomass or zooplankton abundance) beyond their natural cycle. The second step consists in obtaining anomalies by subtracting this seasonality from the original time-series. The method used is the seasonal differentiation by the seasonal deviation methods. Finally, the cumulative sum of these anomalies was produced to detect regime shifts in the time-series for the assessment and comparison periods. A Spearman rank correlation test is now implemented to test the trend of the cumulative sum of the anomalies of the assessment and comparison periods. The correlation can move toward a significant ( $p \leq 0,05$ ) increase in phytoplankton biomass/zooplankton abundance (0 to 1), no changes (=0) or decrease in phytoplankton biomass/zooplankton abundance (-1 to 0). The results of the Spearman rank correlation provided an indication of changes. A t-test against the cumulative sum of the anomalies of the comparison period and the assessment period provide information whether the trends are significantly different or not. These improvements to the methodology since IA2017 now allow us to define distinct

comparison and assessment periods. For the QSR2023 assessment, the comparison period included all data prior to 2015 and the assessment period was set from 2015 to 2019, due to post-2019 data not yet being available across all plankton datasets.

### **Spatial scales**

Because plankton community composition, distribution, and dynamics are closely linked to their environment, the analysis was performed at the scale of the 'COMP4 assessment units' (COMP4 v8a; **Figure a, Table a**). Assessment units within the Greater North Sea and Celtic Seas (OSPAR Regions II and III, respectively) were initially developed by Deltares and partner institutes as part of the EU Joint Monitoring Programme of the Eutrophication of the North Sea with Satellite data (JMP-EUNOSAT; Enserink *et al.*, 2019) and further refined in the revision process of the eutrophication assessment by OSPAR expert groups ICG-EMO and TG-COMP. Assessment units with similar phytoplankton dynamics were derived from cluster analysis of satellite data for chlorophyll-*a* and primary production. Boundaries between assessment units were derived by relating clustering results to the best-matching gradients in environmental variables obtained from the three-dimensional hydrodynamic Dutch Continental Shelf model version 6 (DCSMv6 FM). The variables which best matched the divisions highlighted by clustering were depth, salinity, and stratification regime. Additional geographic areas were added such as the Channel, Irish Sea and Kattegat. These assessment units are a geographical representation of the conditions which best suit plankton distribution, dynamics, and community composition.

Because the Bay of Biscay and Iberian Coast (OSPAR Region IV) extended beyond the boundaries of the DCSMv6 FM, assessment units within this Region were developed using a different methodology, based on phytoplankton dynamics (Spain) and salinity dynamics (Portugal). To delineate assessment units for the Spanish coast, a polygon was created to extend from the coast to the exclusive economic zone (EEZ) boundary. Daily MODIS-Aqua Level-2 satellite images were used to calculate climatological mean values of chlorophyll-*a* for each pixel. K-means clustering was then used to group pixels with similar dynamics, resulting in six distinct groupings within the main Spanish polygon. Portugal's three Water Framework Directive assessment units were extended to the boundaries of the Portuguese EEZ. These assessment units were further divided longitudinally to separate pelagic waters from coastal waters more subject to eutrophication from river influence by applying a salinity threshold, followed by a bathymetry threshold.

Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton




Figure a: COMP4 assessment units developed by JMP-EUNOSAT and OSPAR.

**Classification of the pelagic habitats**

Following the European Commission (2017) outlining criteria and methodological standards on good environmental status of marine waters, the COMP4 assessment units and the fixed-point stations are associated with a habitat type within their corresponding OSPAR Region (**Table a**). Habitat identifications were processed following strict criteria according to surface mean salinity and mean depth. Four habitats were identified: variable salinity (corresponding to river plumes and regions of freshwater influence (ROFI)), coastal habitat (nearshore areas adjacent to ROFIs with mean salinity < 34,5), shelf habitat (corresponding to offshore areas with mean depth less than 200 m and mean salinity > 34,5) and oceanic/beyond shelf habitats (corresponding to offshore areas with mean depth greater than 200 m).

**Table a: classification of the COMP4 assessment units and fixed-point stations by habitat type within OSPAR regions.**

Area code	Area name	Salinity (surface mean)	Depth (mean)	Habitat type	OSPAR region
ADPM	Adour plume	34,4	87	Variable salinity	IV
ELPM	Elbe plume	30,8	18		II
EMPM	Ems plume	31,4	19		II
GDPM	Gironde plume	33,5	34		IV
HPM	Humber plume	33,5	16		II
LBPM	Liverpool Bay plume	30,6	15		III
LPM	Loire plume	33,8	38		IV
MPM	Meuse plume	29,3	16		II
RHPM	Rhine plume	31,0	17		II
SCHPM1	Scheldt plume 1	31,4	13		II

SCHPM2	Scheldt plume 2	30,9	15		II
SHPM	Shannon plume	34,1	61		III
SPM	Seine plume	31,8	25		II
THPM	Thames plume	34,4	22		II
CFR	Coastal FR Channel	34,2	33	Coastal	II
CIRL	Coastal IRL 3	34,0	65		III
CNOR1	Coastal NOR 1	34,3	190		II
CNOR2	Coastal NOR 2	34,0	217		II
CNOR3	Coastal NOR 3	32,4	171		II
CUK1	Coastal UK 1	34,5	60		III
CUKC	Coastal UK Channel	34,8	37		II
CWAC	Coastal Waters AC	No information	No information		IV
CWBC	Coastal Waters BC	No information	No information		IV
CWCC	Coastal Waters CC	No information	No information		IV
ECPM1	East Coast (permanently mixed) 1	34,8	73		II
ECPM2	East Coast (permanently mixed) 2	34,5	43		II
GBC	German Bight Central	33,4	39		II
IRS	Irish Sea	33,7	65		III
KC	Kattegat Coastal	25,7	21		II
KD	Kattegat Deep	27,6	50		II
NAAC1A	NorAtlantic Area NOR-NorC1	No information	No information		IV
NAAC1B	NorAtlantic Area NOR-NorC1	No information	No information		IV
NAAC1C	NorAtlantic Area NOR-NorC1	No information	No information		IV
NAAC1D	NorAtlantic Area NOR-NorC1	No information	No information		IV
NAAC2	NorAtlantic Area NOR-NorC2	No information	No information	IV	

Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

NAAC3	NorAtlantic Area NOR-NorC3	No information	No information		IV
OC	Outer Coastal DEDK	33,4	27		II
SAAC1	SudAtlantic Area SUD-C1	No information	No information		IV
SAAC2	SudAtlantic Area SUD-C2	No information	No information		IV
SAAP2	SudAtlantic Area SUD-P2	No information	No information		IV
SNS	Southern North Sea	34,3	32		II
ASS	Atlantic Seasonally Stratified	35,2	134	Shelf	III, IV
CCTI	Channel Coastal shelf tidal influenced	34,8	40		II
CWM	Channel well mixed	35,1	77		II, III
CWMTI	Channel well mixed tidal influenced	35,0	59		II
DB	Dogger Bank	35,1	28		II
ENS	Eastern North Sea	34,8	43		II
GBCW	Gulf of Biscay coastal waters	34,6	53		IV
GBSW	Gulf of Biscay shelf waters	34,9	107		IV
IS1	Intermittently stratified 1	35,3	138		II, III
IS2	Intermittently stratified 2	35,1	102		II
NAAP2	NorAtlantic Area NOR-NorP2	No information	No information		IV
NAAPF	NorAtlantic Area NOR-Plataforma	No information	No information		IV
NNS	Northern North Sea	35,0	121		II
NT	Norwegian Trench	34,1	349		II
SAAP1	SudAtlantic Area SUD-P1	No information	No information		IV
SK	Skagerrak	31,8	134		II
SS	Scottish Sea	35,1	89		II, III
ATL	Atlantic	35,3	2291		II, IV, V

NAAO1	NorAtlantic Area NOR-NorO1	No information	No information	Oceanic / Beyond shelf	IV
OWAO	Ocean Waters AO	No information	No information		IV
OWBO	Ocean Waters BO	No information	No information		IV
OWCO	Ocean Waters CO	No information	No information		IV
SAAOC	Sudatlantic Area SUD-OCEAN	No information	No information		IV
Stonehaven		No information	No information	Coastal	II
Loch Ewe		No information	No information	Variable salinity	III
L4		No information	No information	Coastal	II
N14 Falkenberg		No information	No information	Coastal	II
Anholt		No information	No information	Coastal	II
Slaggö		No information	No information	Coastal	II
E2CO		No information	No information	Coastal	IV
E3VI		No information	No information	Coastal	IV
A17		No information	No information	Shelf	II
E1GI		No information	No information	Shelf	IV
E2GI		No information	No information	Shelf	IV
E3GI		No information	No information	Shelf	IV
E1CU		No information	No information	Shelf	IV
E2CU		No information	No information	Shelf	IV
E3CU		No information	No information	Shelf	IV

#### Data provided and used in this assessment

Satellite data for the Greater North Sea (OSPAR Region II) and the Celtic Seas (OSPAR Region III) were provided by the Royal Belgian Institute for Natural Sciences and Plymouth Marine Laboratory. Missing values or gaps smaller than 4 months were interpolated by cubic spline method. In addition to satellite data, *in situ* chlorophyll-*a* data from stations were provided by Plymouth Marine Laboratory for L4 station (Channel well mixed assessment unit), by Marine Scotland Science for Stonehaven and Loch Ewe fixed stations, by the Swedish Meteorological and Hydrological Institute and by Aarhus University (Kattegat and Skagerrak assessment units) and by Landesamt für Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein and the Niedersächsischer Landesbetrieb für Wasserwirtschaft, Küsten und Naturschutz (North Sea). In the Bay of Biscay and Iberian Coast (OSPAR Region IV), data were provided by the Instituto Espanol de Oceanografia from a set of fixed stations located in the Bay of Biscay and Iberian Coast.

In addition, non-station data from the CPR have been provided respectively by the Marine Biological Association. CPR data were used for calculating the PH2 indicator on zooplankton (copepod) abundance. However, the CPR Phytoplankton Colour Index (PCI) was not used in the end for phytoplankton biomass assessment, despite CPR providing the longer time series that would be possible to compare to phytoplankton time-series. For phytoplankton biomass, it is more suitable to use satellite data as they provide a monthly integrated and synoptic view of chlorophyll-*a* across the Greater North Sea and the Celtic Seas (OSPAR Regions II and III respectively).

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

In a few cases, the spatially distributed data resulted in localised groupings of samples being extrapolated across much larger assessment units (see distribution of CPR samples in **Figure d (i)** ([PH1 indicator assessment](#)), as was the case for CPR data along the west of Scotland (Intermittently Stratified 1) and Ireland (Atlantic Seasonally Stratified).

Additional data from fixed stations were provided by Belgium, the Netherlands, and the United Kingdom, however, due to some time-series being shorter than the comparison-assessment period or located outside OSPAR Regions, some data were not used in this assessment. Data from ongoing monitoring within OSPAR assessment regions will be used in the future (e.g., VLIZ data from Belgium). **Table b** provides a list of the data used for this assessment.

**Table b: Contracting Parties and institutes that provided the datasets used for this pelagic assessment.**

Contracting Party	Institute	Dataset name	Date range	Sampling frequency
Belgium	Royal Belgium Institute of Natural Sciences (RBINS)	CHL_RBINS	2009-2020	Daily
Denmark	Aarhus University	NOVANA chlorophyl data	2009-2020	Fortnightly
Germany	Landesamt für Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein (LLUR)	OSPAR_LLUR-SH Phytoplankton Biomass_2010-2020	2010-2020	Fortnightly
	Niedersächsischer Landesbetrieb für Wasserwirtschaft, Küsten und Naturschutz (NLWKN)	OSPAR_NLWKN_1999-2019	1999-2019	Fortnightly to higher frequency
Spain	Instituto Espanol de Oceanografia (IEO)	IEO_RADIALES_Chla	1989-2020	Monthly
		IEO_RADIALES_Zoo	1991-2018	Monthly
Sweden	Swedish Meteorological and Hydrological Institute (SMHI)	National data_SMHI phytoplankton biomass	1986-2020	Monthly
		National data_SMHI_zoo	1996-2020	Monthly
United Kingdom	Environment Agency (EA)	EA CHL 2000-2020	2000-2020	Weekly or higher frequency
	Marine Biological Association (MBA)	CPR dataset 1960-2019	1960-2019	Weekly or higher frequency
	Marine Scotland Science (MSS)	MSS Loch Ewe biomass	2002-2020	Weekly
		MSS Loch Ewe zooplankton	2002-2017	Weekly
		MSS Stonehaven biomass	1997-2020	Weekly
		MSS Stonehaven zooplankton	1999-2020	Weekly
	Plymouth Marine Laboratory (PML)	PML_L4 chl a	1992-2020	
		PML_L4 zooplankton	1988-2020	Fortnightly to higher frequency
PML ICES satellite		1997-2016	Monthly	

### Relationship between environmental pressures and phytoplankton biomass / zooplankton abundance

Environmental variables were selected according to their relevance and availability for plankton to determine the most important pressure in phytoplankton biomass and zooplankton abundance changes. While all the variables listed in **Table c** were used for modelling phytoplankton biomass, nutrients were removed from the model explaining zooplankton abundance. The set of environmental variables used originated from different models targeting the North-East Atlantic area.

The first step consisted of evaluating long-term links to pressures and to avoid excluding the first several decades of many plankton time-series due to missing values. To achieve this, the method used multiple random forest regressions to impute missing values based on collinearities among observed values in the predictors. For each variable containing missing values the algorithm generated a separate regression model based on all the other predictors. To improve imputation performance, a numeric variable representing 'month' was included in this step to better predict the consistent seasonal patterns in some variables. This step was performed using 'missRanger' R package (Mayer and Mayer 2019).

Then, values for each environmental variable were calculated as the mean of monthly mean gridded values (modelled and remotely sensed) within each COMP4 assessment unit. For fixed-point stations, mean values were calculated from all measurements within a 5-nautical mile radius of the station. Where *in situ* data were available (total nitrogen, nitrate, phosphate, total phosphorous, silicate) they were evaluated instead of the modelled environmental variables. For Atlantic Multidecadal Oscillation (AMO) and North Atlantic Oscillation (NAO), monthly values were applied identically across all assessment units since these variables have basin-scale influence likely to cover the entire assessment region.

Finally, a random forest algorithm was applied to evaluate which was the best combination of environmental variable for phytoplankton biomass / zooplankton abundance prediction. The algorithm is based on the combination of predictions made by multiple regression trees (here,  $k = 1000$  trees) with the optimal tree (defined as the best combination of variables) obtained by majority voting. It is however important to note that the random forest algorithm may be affected by correlated variables.

Prior to analysis, the original datasets were split into two subsets resulting in a training set and a test set. The training set was used for the selection of the best combination while the test set was used to validate the predictions. The training set consisted of data from the comparison period (prior to 2015) while the test set consisted of data from the assessment period (from 2015 to 2019).

For map visualisation, some variables were aggregated together. Total nitrogen, nitrate, phosphate, total phosphorus, N:P ratio and silicate were pooled under the term "nutrient". The same procedure was applied for AMO and NAO which were pooled under the term "Climate indices".

It is important to keep in mind that on the pressures presented below, pH is not an independent variable. Indeed, the pH of seawater is controlled by the concentration of dissolved inorganic carbon (DIC). Changes in pH can be linked to changes in phytoplankton activity, since phytoplankton ingest DIC to fuel growth and reproduction, but also to the concentration of atmospheric carbon dioxide, the main driver of global warming. Consequently, increases of phytoplankton biomass could increase seawater pH and vice versa. However, a change in pH can adversely affect phytoplankton biomass.

**Table c: list of environmental variables used as pressures.**

Variable name	Description	Abbreviation	Source
Sea surface	Temperature of surface	sst	International Comprehensive Ocean-Atmosphere Data Set (ICOADS): <a href="https://psl.noaa.gov/data/gridded/data.coads.1deg.html">https://psl.noaa.gov/data/gridded/data.coads.1deg.html</a>

Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

temperature	layer, as measured by satellite		
Salinity	Salinity of the surface layer	sal	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_PHY_004_009; <a href="https://doi.org/10.48670/moi-00059">https://doi.org/10.48670/moi-00059</a> )
Total Oxidised Nitrogen	Total oxidised nitrogen concentration in surface layer	totn	In situ data from Marine Scotland Science (MSS): <a href="https://doi.org/10.7489/1881-1">https://doi.org/10.7489/1881-1</a>
Nitrate	Nitrate concentration of the surface layer	ntra	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_BGC_004_011): <a href="https://doi.org/10.48670/moi-00058">https://doi.org/10.48670/moi-00058</a> ; In situ data from Plymouth Marine Laboratory (PML): <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a>
Phosphate	Dissolved inorganic phosphate concentration of the surface layer	phos	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_BGC_004_011): <a href="https://doi.org/10.48670/moi-00058">https://doi.org/10.48670/moi-00058</a> ); In situ data from Plymouth Marine Laboratory (PML): <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a> and Marine Scotland Science (MSS): <a href="https://doi.org/10.7489/1881-1">https://doi.org/10.7489/1881-1</a> <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a>
Total phosphorous	Total phosphorous concentration of the surface layer	totp	In situ data from Aarhus University (Svendsen <i>et al.</i> 2005)
N:P ratio	The ratio of molar nitrogen concentration to molar phosphorus concentration	np	Derived from: European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_BGC_004_011): <a href="https://doi.org/10.48670/moi-00058">https://doi.org/10.48670/moi-00058</a> ; In situ data from Plymouth Marine Laboratory (PML): <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a> and Marine Scotland Science (MSS): <a href="https://doi.org/10.7489/1881-1">https://doi.org/10.7489/1881-1</a>
Silicate	Dissolved	Si	In situ data from Marine Scotland Science (MSS): <a href="https://doi.org/10.7489/1881-1">https://doi.org/10.7489/1881-1</a> ;

	silicate concentration of the surface layer		Plymouth Marine Laboratory (PML): <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a> ; In situ data from Aarhus University (Svendsen <i>et al.</i> 2005)
Wind speed	Wind speed (proxy of turbulence)	wspd	International Comprehensive Ocean-Atmosphere Data Set (ICOADS); <a href="https://psl.noaa.gov/data/gridded/data.coads.1deg.html">https://psl.noaa.gov/data/gridded/data.coads.1deg.html</a>
Mixed layer depth	Surface layer in which density is nearly homogeneous with depth	mld	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_PHY_004_009); <a href="https://doi.org/10.48670/moi-00059">https://doi.org/10.48670/moi-00059</a>
Light attenuation	The extinction coefficient for the visible light in the water column	attn	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_BGC_004_011); <a href="https://doi.org/10.48670/moi-00058">https://doi.org/10.48670/moi-00058</a> <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a> <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a>
Precipitation	Rate of precipitation	precip	International Comprehensive Ocean-Atmosphere Data Set (ICOADS); <a href="https://psl.noaa.gov/data/gridded/data.coads.1deg.html">https://psl.noaa.gov/data/gridded/data.coads.1deg.html</a>
Current velocity	Current velocity in the surface layer	cvel	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_PHY_004_009); <a href="https://doi.org/10.48670/moi-00059">https://doi.org/10.48670/moi-00059</a>
pH	Sea water pH reported on total scale	pH	European Regional Seas Ecosystem Model (ERSEM; NWSHELF_MULTIYEAR_BGC_004_011); <a href="https://doi.org/10.48670/moi-00058">https://doi.org/10.48670/moi-00058</a> <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a> <a href="https://www.westernchannelobservatory.org.uk/l4_nutrients.php">https://www.westernchannelobservatory.org.uk/l4_nutrients.php</a>
NAO	The North Atlantic Oscillation	nao	National Oceanic and Atmospheric Administration (NOAA): <a href="https://www.ncdc.noaa.gov/teleconnections/nao/">https://www.ncdc.noaa.gov/teleconnections/nao/</a>

Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

	on is a weather phenomenon over the North Atlantic Ocean of fluctuations in the difference of atmospheric pressure at sea level between the Icelandic Low and the Azores High		
AMO	The Atlantic Multidecadal Oscillation is the theorised variability of the sea surface temperature of the North Atlantic Ocean on the timescale of several decades	amo	National Oceanic and Atmospheric Administration (NOAA); <a href="https://psl.noaa.gov/data/timeseries/AMO/">https://psl.noaa.gov/data/timeseries/AMO/</a>

**Integration of indicator results:**

A primary objective of this indicator assessment was to integrate results to facilitate an understanding of changes occurring across pelagic habitat types within OSPAR Regions II, III and IV. This required indicator results for each of these OSPAR Regions to be integrated according to the following pelagic habitat

categories: variable salinity, coastal, shelf, and oceanic / beyond shelf. This categorisation of COMP4 assessment units and fixed-point stations is described in **Table a**. To meet this objective, we focused on the primary direction of change detected across assessment units and fixed-point stations within each pelagic habitat category for each of the 2 plankton components highlighted in this assessment. We then reported the mean confidence, spatial representativeness, and most likely links to environmental pressures.

As an example, changes in zooplankton were assessed across 6 COMP4 areas and 4 fixed-point stations representing coastal habitats within OSPAR Region II. If 4 decreasing trends, 3 increasing trends, and 3 instances of no trend were detected across these locations, we would report a decreasing net trend and the proportion of assessment units studied where this trend was detected, in this case 0,4.

The spatial representativeness of the result corresponds the proportion of the total number of COMP4 assessment units considered in the analysis, in this case 6, out of the total number of possible COMP4 assessment units representing coastal habitats within the OSPAR Region, in this case 12. Therefore, the spatial representativeness of the result would be 0,5. Note that fixed-point station datasets do not contribute to this score.

Finally, to report links to environmental pressures which can drive changes in PH2 components for the net trend, we ranked environmental variables for each location based on their relative variable importance, with 1 assigned to the variable with highest importance, 2 to the variable with second highest importance and so on. For locations where the net trend was increasing, we calculated the mean rank of each environmental variable and reported the variable with the lowest mean rank.

**Addressing PH2 quality status**


In order to deliver a clear and comprehensive message to the scientific and non-scientific community, the results of the indicator were summarised by their quality status per habitat within each OSPAR region. The quality status had been defined by the change in indicator value for each plankton component according to assessment threshold and / or the impact of anthropogenic pressures and climate change on the indicator change (McQuatters-Gollop *et al.*, 2022). Thus, the quality status resulted in 4 categories: Not good, Unknown, Good, and Not assessed. **Table d** gives the detailed explanation of the different categories.

**Table d:** Categorization of the quality status and their associated narratives.


Quality status categories	
Not good	Indicator value is below assessment threshold, or change in indicator represents a declining state, or indicator change is linked to increasing impact of anthropogenic pressures (including climate change), or indicator shows no change but state is considered unsatisfactory
Unknown	No assessment threshold and/or unclear if change represents declining or improving state, or indicator shows no change but uncertain if state represented is satisfactory
Good	Indicator value is above assessment threshold, or indicator represents improving state, or indicator shows no change but state is satisfactory
Not assessed	Indicator was not assessed in a region due to lack of data, lack of expert resource, or lack of policy support.

## Résultats

Cette évaluation identifie les modifications de la biomasse du phytoplancton et de l'abondance du zooplancton à l'échelle régionale et locale. Parmi les 64 zones COMP4, des changements significatifs ont été constatés dans 39 unités d'évaluation pour la biomasse du phytoplancton (**Figure 1**) et dans 22 unités d'évaluation pour l'abondance du zooplancton (**Figure 2**).



**Figure 1 : Tendance des anomalies de la biomasse phytoplanctonique entre la période d'évaluation (2015-2019) et la période de comparaison (données des stations : 1992-2014 ; données des non-stations : 1997-2014). Les zones hachurées sont caractérisées par des changements significatifs ( $p \leq 0,05$ ) de la biomasse de phytoplancton entre la période de comparaison et la période d'évaluation. Les zones blanches indiquent l'absence de données ou des données insuffisantes pour évaluer la zone.**




**Figure 2 : Tendence des anomalies de l'abondance zooplanctonique entre la période d'évaluation (2015-2019) et la période de comparaison 1960-2014). Les zones hachurées sont caractérisées par des changements significatifs ( $p \leq 0,05$ ) de l'abondance zooplanctonique entre la période de comparaison et la période d'évaluation. Les zones blanches indiquent l'absence de données ou des données insuffisantes pour évaluer la zone. Les résultats des unités d'évaluation de la stratification intermittente 1 (Écosse) et de la stratification saisonnière de l'Atlantique (Irlande) doivent être interprétés avec prudence, car les échantillons du CPR ont été extrapolés sur des unités d'évaluation beaucoup plus vastes.**

Les anomalies de la biomasse du phytoplancton et de l'abondance du zooplancton ont présenté des tendances à la baisse dans la majorité des unités COMP4 évaluées. La période d'évaluation (2015-2019) a été marquée par des diminutions fortes et significatives de la biomasse de phytoplancton dans 32 unités d'évaluation, tandis que des augmentations ont été documentées dans 7 zones. En ce qui concerne l'abondance du zooplancton, de fortes diminutions ont eu lieu dans 13 zones, tandis que l'abondance a augmenté dans 8 zones et qu'une zone n'a connu aucun changement. La biomasse du phytoplancton a fortement diminué dans les eaux du large (zones atlantiques et golfe de Gascogne), dans la Manche et dans quelques zones de la mer du Nord (principalement la mer du Nord orientale et la fosse norvégienne), tandis qu'elle a augmenté dans une grande partie des zones de la mer du Nord (panaches de l'Elbe, de l'Ems et de l'Humber, mer du Nord septentrionale et méridionale, banc du Dogger). En ce qui concerne le zooplancton, une forte diminution de l'abondance a été observée dans les eaux du large telles que les eaux de haute mer (zones atlantiques et golfe de Gascogne), la Manche et une grande partie de la mer du Nord (à l'exception des eaux côtières et de la fosse norvégienne). Une augmentation significative de la biomasse phytoplanctonique a été observée dans les habitats côtiers et les habitats du plateau de la Mer du Nord au sens large et des Mers celtiques (canal côtier du Royaume-Uni et canal côtier du Royaume-Uni 1, mers d'Irlande et d'Écosse, côte est en permanence mixte 1, fosse norvégienne).


## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

En accord avec les résultats de cet indicateur présentés dans l'IA2017, la biomasse phytoplanctonique a continué à augmenter en mer du Nord, sauf dans la partie orientale de la mer du Nord (mer du Nord orientale, German Bight Central and Outer Coastal DEDK). L'abondance du zooplancton a fortement diminué dans les stations des eaux de haute mer (zones atlantiques et golfe de Gascogne), dans certaines régions de la Manche (Manche bien mélangée et Manche bien mélangée influencée par les marées) et dans les eaux de la mer du Nord (mer du Nord septentrionale, Dogger Bank, mer du Nord orientale, German Bight Central et Outer Coastal DEDK). Une augmentation de l'abondance a été principalement observée le long des côtes anglaises et dans la fosse norvégienne. Bien que l'IA2017 ait signalé une tendance négative dans l'abondance du zooplancton pour le sud de la mer du Nord, la période d'évaluation actuelle (2015-2019) n'a détecté aucun changement significatif dans l'abondance du zooplancton pour cette zone. Les résultats peuvent être sujets à des variations par rapport à la littérature existante, probablement en raison de différences dans l'agrégation spatiale avant l'analyse.

Des liens entre la biomasse du phytoplancton (**Figure 3**) et l'abondance du zooplancton (**Figure 4**) et le contrôle des pressions liées au changement climatique ont été mis en évidence dans les mers celtiques, le golfe de Gascogne et la côte ibérique. L'augmentation de la température de surface de la mer (SST), la diminution de la vitesse du vent, la diminution de l'atténuation de la lumière et l'augmentation de la profondeur de la couche mélangée ont été liées à la diminution de la biomasse du phytoplancton et de l'abondance du zooplancton. En outre, les diminutions de la biomasse du phytoplancton et de l'abondance du zooplancton dans les habitats côtiers et du plateau des Mers celtiques ont été associées à la baisse du pH. Dans les habitats côtiers et de plateau à salinité variable de la grande mer du Nord, les nutriments (nitrate, phosphate, rapport N:P), considérés comme un indicateur de l'eutrophisation, ont été les variables les plus importantes dans les changements de la biomasse du phytoplancton. L'impact de l'eutrophisation peut être significatif dans cette région et des investigations supplémentaires doivent être menées avec le groupe GIC-Eut d'OSPAR afin de mieux comprendre les relations sous-jacentes. En ce qui concerne ces relations, l'état de qualité de la plupart des habitats dans les Régions OSPAR est « Pas bon ». Seuls les habitats à salinité variable de la Mer du Nord au sens large et des Mers celtiques ont un état de qualité "Inconnu". Des détails supplémentaires pour cette section se trouvent dans la section « résultats étendus ».



**Figure 3 : Variables les plus importantes concernant les modifications de la biomasse du phytoplancton dans les zones d'évaluation du programme COMP4. Disponible via : [ODIMS](#)**




**Figure 4 : Variables les plus importantes concernant les modifications de l'abondance du zooplancton dans les zones d'évaluation du programme COMP4. Disponible via : [ODIMS](#)**

## Résultats (version étendue)


### Anomalies percentile in the time-series

For the previous assessment (IA2017) annual anomalies in a time-series were categorised as either small, intermediate, or important changes. This categorisation could be used as an “early warning” signal of change. Some examples of anomalies from the Dogger Bank are displayed for phytoplankton biomass (**Figure b**) and for zooplankton abundance (**Figure c**) using the previous assessment methods based on percentiles. This method focused on the amplitude of monthly deviation from mean conditions, rather than assessing a trend through time. For example, extreme values [ $<2,5$  or  $>97,5$ ] accompanied large changes in the plankton, triggering further investigation into understanding such changes.

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton




**Figure b: Monthly anomalies of phytoplankton biomass (non-station data from satellite) for the Dogger Bank area over the period 1997-2019. Three colours have been attributed to each category: Light blue (small change), blue (intermediate change) and dark blue (important change).**



**Figure c: Monthly anomalies of zooplankton abundance (non-station data from the CPR) for the Dogger Bank area over the period 1960-2019. Three colours have been attributed to each category: Light blue (small change), blue (intermediate change) and dark blue (important change).**

### Integration of phytoplankton biomass and zooplankton abundance trends

The results of the current assessment have indicated 3 patterns of change between phytoplankton and zooplankton (**Figure d**): **(i)** a simultaneous decrease in both phytoplankton biomass and zooplankton abundance, **(ii)** increase in phytoplankton biomass and decrease in zooplankton abundance or decrease in phytoplankton biomass and increase in zooplankton abundance, and **(iii)** a simultaneous increase of phytoplankton biomass and zooplankton abundance.



**Figure d: Integration between PH2 phytoplankton and zooplankton trends within the different COMP4 assessment units and at the different fixed stations. Downward arrow represents simultaneous decrease of phytoplankton biomass and zooplankton abundance (pattern i). Upward arrow represents simultaneous increase of phytoplankton biomass and zooplankton abundance (pattern iii). Bidirectional arrows represent increase of one plankton component while the second component decreases and vice versa (pattern ii).**

The first pattern **(i)** was observed in the open ocean waters (Atlantic areas), as well as in the Channel well mixed and Channel well mixed tidal influenced areas, in the south-eastern part of the North Sea (e.g., Eastern North Sea, German Bight Central) as well as in some fixed-points (i.e. L4, Loch Ewe, E1GI, E2GI, Anholt and Slaggö).

The second pattern **(ii)** was recorded in the Northern North Sea, the Dogger Bank, in the Coastal Waters of Ireland, Outer Coastal waters, along the Southern English coasts (Coastal waters of Southern England), the Irish and Scottish Seas, and in the North Sea (East Coast of England, Intermittently stratified waters, Coastal waters of Norway and the Norwegian Trench). This pattern was found in only one monitoring station: Stonehaven location.

The third pattern **(iii)** was observed in the Elbe and Ems plumes and in the fixed points in the Skagerrak (i.e., Slaggö, A17) and the Iberian coast (i.e., E1CU, E2CU, E3CU, E3GI and E2CO).

Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

### **Relation between pressures and indicator**

In the Greater North Sea, modelling results have revealed nutrients (dissolved inorganic phosphate, nitrogen, and the balance between nitrogen and dissolved inorganic phosphate) as the most important pressures affecting changes in phytoplankton biomass for the three types of habitats (variable salinity, coastal and shelf). Variable salinity habitat was marked by a decrease in phosphate which in turn was high correlated with a decrease of phytoplankton biomass (see also Gohin *et al.*, 2020). Faster reduction of phosphorous than nitrogen occurred in both coastal and shelf habitats resulting in increasing N:P ratio. In both habitats, these conditions occurred while phytoplankton biomass decreased. Zooplankton abundance was rather linked to direct or indirect climatic conditions. Despite increasing, SST were reported as the most important variable for zooplankton abundance decrease in the variable salinity habitat, the relation was insignificant thus changes did not occur during the assessment period. Finally, decreasing mixed layer depth (MLD) was linked with decreasing zooplankton abundance in coastal habitat, decreasing MLD was linked with increasing zooplankton abundance in shelf habitats.

In the Celtic Seas, the current assessment showed that decreasing pH was most likely linked with phytoplankton biomass decrease in coastal habitat and in shelf habitats for both phytoplankton biomass and zooplankton abundance (further explanations on the relationship between pH and phytoplankton biomass are given in the subsection Relationship between environmental pressures and phytoplankton biomass / zooplankton abundance). Changes in pH can be linked to changes in phytoplankton activity, since phytoplankton ingest dissolved inorganic carbon (DIC) to fuel growth and reproduction, but also to the concentration of atmospheric carbon dioxide, the main driver of global warming. However, for reasons described above, the relationship between pH and phytoplankton biomass should be evaluated cautiously and further analysis is necessary to quantify phytoplankton's contribution to pH variability. Indicator results in variable salinity habitats also reflected a reduction in phytoplankton biomass due to decreasing nitrate concentration. The increase in zooplankton abundance in the coastal habitat was linked to natural climatic condition. Finally, decrease in zooplankton abundance in variable salinity habitats was linked to decrease of current velocity.

In the Bay of Biscay and Iberian Coast (OSPAR Region IV), changes in both phytoplankton biomass and zooplankton abundance were most strongly linked to climatic conditions. While phytoplankton biomass decreased in shelf habitats due to increasing mixed layer depth over the last decades, phytoplankton biomass also decreased in oceanic habitats, associated with a decrease in light attenuation and thus shallowing of the euphotic zone. Surprisingly, zooplankton abundance in coastal and oceanic habitats was positively correlated with SST. Results indicated that abundance increased as well as SST in coastal habitat whereas abundance and SST both decreased in oceanic habitats. Finally, decreases in zooplankton abundance in shelf habitats of the Bay of Biscay and Iberian Coast were linked to increasing mixed layer depth.

The quality status of the pelagic habitats within OSPAR Regions was addressed following the links between pressures and PH2 indicator results (**Table e**). Despite relationships with environmental variables remaining unclear, climate change and decrease in pH were linked with the indicators within coastal and shelf habitats of the Celtic Seas and within coastal, shelf and Oceanic / beyond shelf habitats of the Bay of Biscay and Iberian Coast. Consequently, the quality status within these habitats is "Not good". The faster reduction in phosphorus compared to nitrogen has increased the N:P ratio which in turns impacted the PH2 indicator. This was the case for coastal and shelf habitats of OSPAR Region II. As a consequence of the imbalance between nitrogen and phosphorus, this might be related to eutrophication, and the quality status of the habitat in which this observation was made was also determined to be "Not good". Finally, where other parameters were linked with changes in phytoplankton biomass and zooplankton abundance (e.g., variable

salinity habitats of the Greater North Sea and the Celtic Seas), it was more difficult to establish the origin of the pressure. Quality status of the habitat was therefore categorised as “Uncertain”.

**Table e: Integration of the indicator results for the Greater North Sea, the Celtic Seas and the Bay of Biscay and Iberian Coast. Column names are described as follows: Dir: the net direction of change in the plankton component (upward arrow: increasing trend, equality sign: no trend, downward arrow: decreasing trend), Trend: the percentage of assessment units exhibiting the respective trend (if no results were reported for assessment units, stations are used), Change: a logical variable (TRUE/FALSE) to report whether a net trend is likely given the significance of the results, Pressure: the environmental pressure with the greatest mean rank for the respective trend, Rank: the mean rank of the environmental pressure indicated under Pressure, nSt: the total number of monitoring fixed stations considered, nCOMP4: The total number of COMP4 assessment units considered, totCOMP4: The total number of potential COMP4 assessment units for the habitat category, spatialRep: the spatial representativeness score of the analysis.**

OSPAR Region	Habitat	Plankton component	Dir	Trend	Change	Pressure	Rank	nSt	nCOMP4	totCOMP4	Spatial Rep	
The Greater North Sea	Variable salinity	Phytoplankton	↓	60%	TRUE	phosp	2.8	1	9	9	100%	
		Zooplankton	↓	100%	FALSE	sst	1.0	1	0	9	0%	
	Coastal	Phytoplankton	↓	92%	TRUE	np	4.0	1	12	12	100%	
		Zooplankton	↑	40%	TRUE	mld	2.8	4	6	12	50%	
	Shelf	Phytoplankton	↓	75%	TRUE	np	2.3	1	11	11	100%	
		Zooplankton	↓	67%	TRUE	mld	2.7	0	7	11	64%	
	Oceanic	Phytoplankton	NA									
		Zooplankton	NA									
	The Celtic Seas	Variable salinity	Phytoplankton	↓	100%	TRUE	ntra	2.0	1	2	2	100%
			Zooplankton	↓	100%	TRUE	cvel	1.0	1	0	2	0%
Coastal		Phytoplankton	↓	67%	TRUE	pH	1.7	0	3	3	100%	
		Zooplankton	↑	67%	TRUE	AMO	2.0	0	3	3	100%	
Shelf		Phytoplankton	↓	100%	TRUE	pH	3.5	0	4	4	100%	
		Zooplankton	↓	75%	TRUE	pH	2.5	0	4	4	100%	
The Bay of Biscay and Iberian Coast	Oceanic	Phytoplankton	NA									
		Zooplankton	NA									
	Variable salinity	Phytoplankton	NA									
		Zooplankton	NA									
	Coastal	Phytoplankton	↓	100%	TRUE	NAO	1.0	1	0	12	0%	
		Zooplankton	↑	100%	TRUE	sst	2.5	2	0	12	0%	
Shelf	Phytoplankton	↑	43%	TRUE	wspd	1.0	6	1	6	17%		
	Zooplankton	↓	100%	TRUE	mld	3.0	0	2	6	33%		
Oceanic	Phytoplankton	↓	100%	TRUE	attn	1.0	0	1	6	17%		

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

	Zooplankton	↓	100%	TRUE	sst	1.5	0	2	6	33%
--	-------------	---	------	------	-----	-----	---	---	---	-----

### Conclusion

Cet indicateur identifie les modifications de la biomasse du phytoplancton et de l'abondance du zooplancton à l'échelle régionale (unités d'évaluation COMP4) et locale (stations de surveillance discrètes). Les résultats mettent en évidence une diminution générale de la biomasse du phytoplancton et de l'abondance du zooplancton à l'échelle régionale. Toutefois, à l'échelle locale, la biomasse de phytoplancton observée a augmenté depuis l'évaluation précédente en mer du Nord et l'abondance du zooplancton a également augmenté le long des côtes du Royaume-Uni et dans la fosse norvégienne. Des liens possibles avec le changement climatique ont été signalés, en particulier dans les habitats côtiers, les plateaux et les habitats océaniques de la Mer du Nord au sens large et des Mers celtiques. Les nutriments ont été associés à l'indicateur, en particulier dans la Mer du Nord au sens large. De futures études sont nécessaires pour explorer les liens avec les indicateurs pertinents relatifs à l'habitat pélagique, au réseau trophique et à l'eutrophisation.

### Conclusion (version étendue)

This assessment identifies changes in phytoplankton biomass and zooplankton abundance at regional (COMP4 assessment units) and local (discrete monitoring stations) scales. The results highlight a general decrease in phytoplankton biomass and zooplankton abundance at the regional scale. However, at the local scale, observed phytoplankton biomass has been increasing since the previous assessment in the North Sea and zooplankton abundance has also been increasing along the coasts of the United Kingdom and in the Norwegian trench. Possible links with climate change were reported especially in coastal, shelf and oceanic habitats of the Greater North Sea and the Celtic Seas. Eutrophication has been linked to the indicator especially in the Greater North Sea. Future investigations are needed to explore connections with relevant Pelagic Habitat and Food Web indicators.

### Lacunes dans les connaissances

Les travaux supplémentaires recommandés sont: 1. Inclusion d'ensembles de données supplémentaires, en particulier pour le golfe de Gascogne et la côte ibérique ; 2. comparaison avec les indicateurs pertinents de l'habitat pélagique et du réseau trophique afin de mieux évaluer l'ampleur du changement ; 3. poursuite de l'amélioration de la méthodologie afin de définir plus précisément le cycle naturel de la biomasse du phytoplancton et de l'abondance du zooplancton ; 4. affinement de l'utilisation des données satellitaires et amélioration de la cohérence avec les évaluations de l'eutrophisation ; et 5. affinement des liens entre PH2 et les pressions. Affiner les liens entre PH2 et les pressions.

### Lacunes dans les connaissances (version étendue)

Further development of this indicator is needed, particularly on the following points:

- Inclusion of additional data sets to improve the confidence of indicator's assessment results: especially for the Bay of Biscay and Iberian Coast. In addition, increased zooplankton monitoring is needed in variable salinity habitats. Finally, there is a need to include additional datasets to allow comparison between the different monitoring strategies.
- Comparison to relevant Pelagic Habitat and Food Web indicators to facilitate of impacts through the food web.
- Improve coherence and integration between the PH2 indicator and relevant indicators within HASEC group.
- Integration of dataset originated from different chl-a methods: continual assessment of characterising the differences in chl-a methodologies is required.

- Improve the methodology for defining the natural cycle and subsequently for the trend characterisation and spatial and temporal confidence of the results.
- Refinement of the use of remote sensing data for improving coherence with the OSPAR eutrophication assessments at different spatial and temporal scales.
- Refine the links between PH2 and pressures to identify the origin of the pressures.

#### **Inclusion of additional data sets:**

Recommendations were made in the IA2017 regarding spatial gaps in the Bay of Biscay and Iberian Coast and they are still relevant for the current assessment. In addition, in the current assessment some habitats within OSPAR Regions were lacking data. In particular, there is a need for: zooplankton data for variable salinity habitats in the Greater North Sea and the Celtic Seas and both phytoplankton and zooplankton data for variable salinity and coastal habitats in the Bay of Biscay and Iberian Coast.

In addition to ensure the perennity of existing time series, future research and monitoring studies should address the various gaps in monitoring data coverage because these will be a limiting factor for the development of the indicators. Semi-automated methods such as *in vivo* multi-spectral fluorometry and flow cytometry (Houliez *et al.*, 2012; Bonato *et al.*, 2016; Thyssen *et al.*, 2015, Louchart *et al.*, 2020), supplemented by automated image acquisition and analysis which allow higher spatial and temporal resolution should also be considered within existing monitoring programmes. In addition, implementation of these innovative techniques on seasonal (see Jouandet *et al.*, 2020) or monthly sampling programs (e.g., Aardema *et al.*, 2019) can help address pelagic community dynamics in offshore waters thus providing information to define the quality status of not assessed areas or assessed areas with low sampling confidence. Furthermore, inclusion of size fractionated chl-a could be integrated to provide a more precise view of phytoplankton biomass. Size fractionated chl-a could also be useful to integrate Pelagic Habitats indicators (this indicator and the indicator '[Changes in plankton lifeforms](#)') since a recommendation to investigate picoplankton and nanoplankton emerged in the previous version of PH1/FW5 indicator.

#### **Comparison to relevant Pelagic Habitat and Food Web Eutrophication indicators:**

Further work is also needed to integrate pelagic habitat indicators (this indicator, and the indicators on '[Changes in Phytoplankton and Zooplankton Communities](#)' and '[Changes in Plankton Diversity](#)') since this will provide greater confidence in the assessment of changes in plankton, and in their integration with indicators of other ecosystem components, in particular food web indicators. PH2 is a bulk indicator, with the highest level of integration among the Pelagic Habitat indicators (McQuatters-Gollop *et al.*, 2017). Whilst PH2 represents the link between the stocks and the fluxes within the pelagic food webs, it does not account for variability in photosynthetic rates and the actual flow of carbon through the system. Further research on the use of primary production (FW2), as a direct indicator of carbon flow through the food web, should be explored.

#### **Improve coherence and integration with relevant eutrophication indicators:**

Being at the bottom of the trophic network, plankton is highly sensitive to their changing environment. In this context, there is a need to improve the linkage with relevant eutrophication indicators from OSPAR's Hazardous Substances and Eutrophication Committee (HASEC) such as the [nutrient inputs indicator](#). Future investigations will help to address the importance of nutrient inputs within the sub-division of the different OSPAR Regions. Furthermore, it is important to establish better coherence between the PH2 indicator and the [chlorophyll-\*a\* concentration indicator](#) across multiple criteria: data collection (similar data should be used to maintain consistency in results), spatial resolution (similar spatial resolution should be maintained across related indicators), and temporal resolution (consistency should be maintained across related indicators in terms of whether they examine year-round time-series, or productive months only).

#### **Integration of dataset originated from different chl-a methods:**

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

As far as future research is concerned, continual assessment of characterising the differences in Chl-*a* methodologies is required (e.g., between fluorometric, spectrophotometric, HPLC and *in situ* fluorometers [which are prone to quenching] plus CPR colour scale index).

### **Methodology improvement:**

The current assessment is built on the assumption that phytoplankton biomass and zooplankton abundance fluctuate over a consistent annual cycle. However, plankton have large year-to-year variability, and this pattern of variability differs across ecosystems and habitats. For example, periodicities of 12 or 6 months or less have been reported. Shifts in the typical annual cycle have also been detected in the past. Future development should therefore focus on investigating the annual cycle for non-stationary datasets (e.g., Winder and Cloern, 2010) and applying this information to de-trend time-series data. Furthermore, it is crucial to investigate the potential change of seasonality over time. Once the annual mean cycle has been removed, the trend analysis can be conducted using linear modelling. Although this methodology is widely accepted, this does not always match with the dataset. Trends can be linear, cubic, quadratic or absent. Consequently, it is necessary to propose an evaluation tool to define more accurately the type of relationship of the de-trended data over time.

### **Refinement of the use of remote sensing data:**

For future assessments, it would be beneficial to examine satellite data at a finer spatial scale, and improve methods for using satellite remote sensing data in turbid and eutrophic water bodies. This will be particularly useful in variable salinity and coastal habitats of the Greater North Sea, which are typically characterised by high turbidity. It is also important to maintain coherence across multiple criteria: data collection (similar data should be used to maintain consistency in results), spatial resolution (similar spatial resolution should be maintained across related indicators, such as PH1/FW5, PH3, FW2 and [chlorophyll-a concentration indicator](#), and temporal resolution (consistency should be maintained across related indicators in terms of whether they examine year-round time-series, or productive months only). It is also important to maintain similar consistency about whether to use temporal interpolation to fill gaps for unsampled months which often occur during winter.

### **Refine the pressure-PH2 relationship:**

The origins of the different pressures are unclear at this stage. Despite evidence from literature, future work should make the distinction between natural versus human induced pressure as well as the origin of natural pressures (e.g., nutrient originating from river run-off or benthic origin or Atlantic flow).

In addition, introduction of a lag into the variable selection is needed to test for delayed effects of environmental pressures on phytoplankton biomass and zooplankton abundance (e.g., the effects of winter nutrient concentrations on phytoplankton biomass during the growing season).

## **Références**

Aardema, H. M., Rijkeboer, M., Lefebvre, A., Veen, A., & Kromkamp, J. C. (2019). High-resolution underway measurements of phytoplankton photosynthesis and abundance as an innovative addition to water quality monitoring programs. *Ocean Science*, 15(5), 1267-1285.

Aubert A., Rombouts I., Artigas F., Budria A., Ostle C., Padegimas B., McQuatters-Gollop A. 2017, "Combining methods and data for a more holistic assessment of the plankton community", as a contribution to the EU Co-financed EcAprHA project (Applying an ecosystem approach to (sub) regional habitat assessments), deliverable No. 1.2.

Batt, R.D., Brock, W.A., Carpenter, S.R., Cole J.J., Pace M.L., Seekell D.A., 2013. Asymmetric response of early warning indicators of phytoplankton transition to and from cycles. *Theor Ecol.*, 6: 285. doi:10.1007/s12080-013-0190-8

Beaugrand G, Luczak C, Edwards M (2009) Rapid biogeographical plankton shifts in the North Atlantic Ocean. *Glob Change Biol* 15:17901790ift

- Berline L., Ibanez F., Grosjean P. 2009. Roadmap for zooplankton time series analysis. SESAME WP1 report. 20p.
- Bonato, S., Christaki, U., Lefebvre, A., Lizon, F., Thyssen, M., and Artigas, L. F. (2015). High spatial variability of phytoplankton assessed by flow cytometry, in a dynamic productive coastal area, in spring: The eastern English Channel. *Estuarine, Coastal and Shelf Science*, 154, 214-223.
- Budria, A., Aubert, A., Rombouts, I., Oslte, C., Atkinson, A., Widdicombe, C., Goberville, E., Artigas., L.F., Johns, D., Padegimas, B., Corcoran, E., McQuatters-Gollop, A. (2017). Cross-linking plankton indicators to better define GES of pelagic habitats. EcApRHA Deliverable WP1.4. 54 pp.
- Burthe, S. J., Henrys, P. A., Mackay, E. B., Spears, B. M., Campbell, R., Carvalho, L., Dudley, B., Gunn, I. D. M., Johns, D. G., Maberly, S. C., May, L., Newell, M. A., Wanless, S., Winfield, I. J., Thackeray, S. J. and Daunt, F. 2016. Do early warning indicators consistently predict nonlinear change in long-term ecological data?. *J Appl Ecol*, 53: 666–676. doi:10.1111/1365-2664.12519
- Buttay L., Miranda A., Casas G., González-Quirós R., Nogueira E. 2015. Long-term and seasonal zooplankton dynamics in the northwest Iberian shelf and its relationship with meteo-climatic and hydrographic variability. *Journal of Plankton Research* 38(1), 106-121
- De Jonge V.N. 2007. Toward the application of ecological concepts in EU coastal water management. *Marine Pollution Bulletin* 55, 407-414.
- Duflos M., Wacquet G., Aubert A., Rombouts I., Mialet B., Devreker D., Lefebvre A., Artigas L.F. 2018. Évaluation de l'état écologique des habitats pélagiques en France métropolitaine. Rapport scientifique pour l'évaluation 2018 au titre du descripteur 1 de la DCSMM. 333 pp.
- Enserink, L., Blauw, A., van der Zande, D., Markager S. (2019). Summary report of the EU project 'Joint monitoring programme of the eutrophication of the North Sea with satellite data' (Ref: DG ENV/MSFD Second Cycle/2016). 21 pp.
- European Commission, 2017. Commission decision (EU) 2017/848 of 17 May 2017 laying down criteria and methodological standards on good environmental status of marine waters and specifications and standardised methods for monitoring and assessment, and repealing Decision 2010/477/EU. *Official Journal of the European Union L*, 125: 43–74.
- Gohin, F., Bryère, P., Lefebvre, A., Sauriau, P. G., Savoye, N., Vantrepotte, V., Bozec, Y., Cariou, T., Conan, P., Coudray, S., Courtay, G., Françoise, S., Goffart, A., Hernández Fariñas, T., Lemoine, M., Piraud, A., Raimbault, P., & Rétho, M. (2020). Satellite and in situ monitoring of Chl-a, Turbidity, and Total Suspended Matter in coastal waters: experience of the year 2017 along the French Coasts. *Journal of Marine Science and Engineering*, 8(9), 665.
- Gowen R.J., McQuatters-Gollop A., Tett P.M., Bresnan E., Castellani C., Cook K., Forster R., Scherer C., Mckinney A. 2011. The Development of UK Pelagic (Plankton) Indicators and Targets for the MSFD, Belfast, 2011.
- Houliez, E., Lizon, F., Thyssen, M., Artigas, L. F., & Schmitt, F. G. (2012). Spectral fluorometric characterization of Haptophyte dynamics using the FluoroProbe: an application in the eastern English Channel for monitoring *Phaeocystis globosa*. *Journal of Plankton Research*, 34(2), 136-151.

## Modifications de la biomasse du phytoplancton et de l'abondance du zooplancton

Ikeda, T. (1985). Metabolic rates of epipelagic marine zooplankton as a function of body mass and temperature. *Marine Biology*, 85(1), 1-11.

Jouandet, M. P., Lebourg, E., Bruaut, M., Dédécker, C., Louchart, A., Gómez, F., Nowaczyck, A., Wacquet, G., Lizon, F., Crouvoisier, M., Cornille, V., Lefebvre, K., Bouchaud, F., Lécuyer, E., Mériaux, X., Cauvin, A., Didry, M., Delabre, J., Duflos, M., and Artigas, L. F. (2020). Résultats des campagnes océanographiques 2017-2019 en Mers Celtiques et Manche Mer du Nord: Test pour la mise en place du Programme de Surveillance DCSMM. 113 pp.

Louchart, A., deBlok, R., Debuschere, E., Gomez, F., Lefebvre, A., Lizon, F., Mortelmans, J., Rijkeboer, M., Schmitt, F.G., Deneudt, K., Veen, A., and Artigas, L.F. (2020). Automated techniques to follow the spatial distribution of *Phaeocystis globosa* and diatoms spring blooms in the Channel and North Sea, Proceedings of the ICHA2018 (18th International Conference on Harmful algae, Nantes 21-26 October 2018), 4 pp.

Mayer, M., & Mayer, M. M. (2019). Package 'missRanger'. *R Package*.

McQuatters-Gollop, A., Edwards, M., Helaouët, P., Johns, D.G., Owens, N.J.P., Raitos, D.E., Schroeder, D., Skinner, J., and Stern, R.F., 2015. The Continuous Plankton Recorder survey: how can long-term phytoplankton datasets deliver Good Environmental Status? *Estuarine, Coastal and Shelf Science*, 162:88-97

McQuatters-Gollop, A., Johns, D. G., Bresnan, E., Skinner, J., Rombouts, I., Stern, R., Aubert, A., Johansen, M., Bedford, J., and Knights, A., (2017). From microscope to management: the critical value of plankton taxonomy to marine policy and biodiversity conservation. *Marine Policy*, 83, 1-10.

McQuatters-Gollop, A., Guérin, L., Arroyo, N.L., Aubert, A., Artigas, L.F., Bedford, J., Corcoran, E., Dierschke, V., Elliott, S.A.M., Geelhoed, S.C.V., Gilles, A., Gonzalez-Irusta, J.M., Haelters, J., Johansen, M., Le Loc'h, F., Lynam, C.P., Niquil, N., Meakins, B., Mitchell, I., Padegimas, B., Pesch, R., Preciado, I., Rombouts, I., Safi, G., Schmitt, P., Schückel, U., Serrano, A., Stebbing, P., De la Torre, A., Vina-Herbon, C. (2022). Assessing the state of marine biodiversity in the Northeast Atlantic. *Ecological Indicators*, 141, 109148.

OSPAR. 2017. Changes in Phytoplankton Biomass and Zooplankton Abundance. <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/>

OSPAR (in prep). OSPAR Agreement 2019-06. Coordinated Environmental Monitoring and Assessment Programme Guidelines for assessing changes in phytoplankton biomass and zooplankton abundance. [www.ospar.org](http://www.ospar.org)

Rombouts I., Beaugrand G., Fizzala X., Gaill F., Greenstreet S.P.R., Lamare S., Le Loc'h F., McQuatters-Gollop A., Mialet B., Niquil N., Percelay J., Renaud F., Rossberg A.G. and Féral J.P., 2013. Food web indicators under the Marine Strategy Framework Directive: from complexity to simplicity? *Ecological Indicators*, 29: 246–254.

Rombouts, I., Beaugrand, G., Ibañez, F., Gasparini, S., Chiba, S., & Legendre, L. 2009. Global latitudinal variations in marine copepod diversity and environmental factors. *Proceedings of the Royal Society B: Biological Sciences*, 276(1670), 3053-3062.

Svendsen, L., L. van der Bijl, S. Boutrup, and B. Norup. 2005. NOVANA. National Monitoring and Assessment Programme for the Aquatic and Terrestrial. and number: NERI Technical Report.

Thyssen, M., Alvain, S., Lefebvre, A., Dessailly, D., Rijkeboer, M., Guiselin, N., Creach, V., and Artigas, L. F. (2015). High-resolution analysis of a North Sea phytoplankton community structure based on in situ flow cytometry observations and potential implication for remote sensing. *Biogeosciences*, 12(13), 4051-4066.

Tilstone, G. H., Land, P. E., Pardo, S., Kerimoglu, O., & Van der Zande, D. (2022). Threshold indicators of primary production in the north-east Atlantic for assessing environmental disturbances using 21 years of satellite ocean colour. *Science of The Total Environment*, 158757.

Turner, J.T. (2004). The importance of small planktonic copepods and their roles in pelagic marine food webs. *Zoological studies* 43: 255-266

Winder, M., and Cloern, J. E. (2010). The annual cycles of phytoplankton biomass. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1555), 3215-3226.

### Métadonnées d'évaluation

Field	Data Type	
<b>Type d'évaluation</b>	Liste	Évaluation de l'indicateur
<b>Résumé des résultats</b>	URL	<a href="https://odims.ospar.org/en/submissions/ospar_plankton_mass_abundance_msfd_2022_06/">https://odims.ospar.org/en/submissions/ospar_plankton_mass_abundance_msfd_2022_06/</a>
<b>Indicateur ODD</b>	Liste	14.2 D'ici à 2020, gérer et protéger durablement les écosystèmes marins et côtiers, notamment en renforçant leur résilience, afin d'éviter les graves conséquences de leur dégradation et prendre des mesures en faveur de leur restauration pour rétablir la santé la productivité des océans
<b>Activité thématique</b>	Liste	Diversité biologique et écosystèmes
<b>Documentation OSPAR pertinente</b>	Texte	OSPAR Agreement 2010-03 The North-East Atlantic Environment Strategy. Strategy of the OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic 2010–2020  OSPAR Agreement 2016-01 OSPAR Coordinated Environmental Monitoring Programme (CEMP)  OSPAR Agreement 2021-01 Strategy of the OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic 2030
<b>Date de publication</b>	Date	2023-06-30
<b>Conditions d'accès et d'utilisation</b>	URL	<a href="https://oap.ospar.org/fr/politique-de-donnees/">https://oap.ospar.org/fr/politique-de-donnees/</a>
<b>Instantané de données</b>	URL	<a href="https://odims.ospar.org/en/submissions/ospar_plankton_changes_snapshot_2022_06/">https://odims.ospar.org/en/submissions/ospar_plankton_changes_snapshot_2022_06/</a>
<b>Résultats des données</b>	Fichier Zip	<a href="https://odims.ospar.org/en/submissions/ospar_plankton_changes_results_2022_06/">https://odims.ospar.org/en/submissions/ospar_plankton_changes_results_2022_06/</a>



COMMISSION  
**OSPAR**

OSPAR Secretariat  
The Aspect  
12 Finsbury Square  
London  
EC2A 1AS  
United Kingdom

t: +44 (0)20 7430 5200  
e: [secretariat@ospar.org](mailto:secretariat@ospar.org)  
[www.ospar.org](http://www.ospar.org)

**Notre vision est celle d'un océan Atlantique Nord-Est propre, sain et biologiquement diversifié, qui soit productif, utilisé de manière durable et résilient au changement climatique et à l'acidification des océans.**

Publication: 1035/2023

© OSPAR Commission, 2023. Permission may be granted by the publishers for the report to be wholly or partly reproduced in publications provided that the source of the extract is clearly indicated.

© Commission OSPAR, 2023. La reproduction de tout ou partie de ce rapport dans une publication peut être autorisée par l'Editeur, sous réserve que l'origine de l'extrait soit clairement mentionnée.